

REDESCRIPTION OF *MEXECHELES VIRGINIENSIS* MALE (ACARIFORMES: CHEYLETIDAE) FROM THE EUROPEAN PART OF RUSSIA

Aleksandra V. Diumina^{1*} and Vladimir V. Abramov²

¹Russian Academy of Sciences, Zoological Institute, St. Petersburg, Russia

²Independent Researcher; Suvorov, Tula Region, Russia

*corresponding author; e-mail: d_alexia@mail.ru

ABSTRACT: The male of *Mexecheles virginensis* (Baker, 1949) (Acariformes; Cheyletidae) found in galleries of scolytine beetles (Coleoptera: Curculionidae) is recorded for the first time in Russia (Tula Region) and redescribed.

KEY WORDS: Acari, Cheyletidae, systematics, *Mexecheles virginensis*.

DOI: 10.21684/0132-8077-2018-26-2-213-217

INTRODUCTION

The family Cheyletidae (Acariformes: Prostigmata) includes over 440 species in 75 genera (Zhang *et al.* 2011). Most members of this family are predators, but some genera are permanent ectoparasites of small mammals and birds (Volgin 1969; Summers and Price 1970; Bochkov and Fain 2001; Bochkov 2004, 2009). The genus *Mexecheles* De Leon, 1962 currently includes ten species represented by free-living predators and those ecologically associated with scolytine beetle galleries (Coleoptera: Curculionidae) (De Leon 1962; Volgin 1969; Smiley and Mooser 1970; Faungarworn and Lekprayoon 2010). The latter work provides a key to females of all presently known species. The currently known *Mexecheles* species, males, were often described for only four species: *Mexecheles votandinii* (Jeffrey, 1975) in Scotland; *M. shiva* (Bochkov et Ochoa, 2005) and *M. virginensis* (Baker, 1949) in India; and *M. thailandensis* Faungarworn et Lekprayoon, 2010 in Thailand (Jeffrey 1975; Fain and Bochkov 2001; Bochkov and Ochoa 2005; Faungarworn and Lekprayoon 2010).

The fauna of cheyletids in the Suvorov District of the Tula Region, Russia was recently studied by Bochkov and Abramov (2016). Among seven cheyletid mites found in that locality, females of *Mexecheles virginensis* were recorded in the territory of Russia for the first time. This species was originally described from females in Virginia, USA (Baker 1949). Males were described much later based on the material from Kashmir, India (Fain and Bochkov 2001). In the present work, we provide a redescription of *M. virginensis* males, based on a newly collected material in the European part of Russia.

MATERIAL AND METHODS

The specimens, used for redescription, were collected near the city of Suvorov (Tula Region,

Russia) by V.V. Abramov in 2016. Extraction from samples was made with Berlese's funnel; mites were stored in tubes with 70% ethanol, and then were mounted in Hoyer's medium. Drawings were made with a Leica microscope, equipped with a camera lucida and DIC optics. The description of idiosomal setation follows that of Grandjean (1939), as adapted for Prostigmata by Kethley (1990). The nomenclature of leg setae follows that of Grandjean (1944). The description follows modern standards proposed in the recent works on *Mexecheles* and related cheyletids (Fain and Bochkov 2001; Bochkov and Ochoa 2005; Faungarworn and Lekprayoon 2010). All measurements are in micrometres (µm). Mite specimens are deposited at the Zoological Institute of the Russian Academy of Sciences (ZISP), St. Petersburg, Russia.

SYSTEMATICS

Family **Cheyletidae Leach, 1815**

Genus ***Mexecheles* De Leon, 1862**

***Mexecheles virginensis* (Baker, 1949)**

(Figs. 1, 2)

Cheyletia virginensis Baker, 1949: 299, pl. 13, Fig. 95–98.

Paracheyletia virginensis, Volgin 1955: 169.

Acarocheyletia virginensis, Smiley and Moser 1970: 229–236.

Mexecheles virginensis, Volgin 1969: 187, Fig. 190–192; Summers and Price 1970: 47, Fig. 39; Fain and Bochkov 2001: 54, Fig. 5.

Material examined. 4 males (AVB 17-1102-69) from galleries of supposedly *Pitiophthorus micrographus* (Linnaeus) (Curculionidae: Scolytinae) under bark of dead fir tree (*Picea* sp.), Russia, Tula Region, Suvorov District, 54°7'20" N, 36°29'47" E,



Fig. 1. *Mexecheles virginiensis*, dorsal view.




Fig. 2. *Mexecheles virginiensis*. A—ventral view of idiosoma and gnathosoma, B—dorsal view of palp, C—D—dorsal view of tarsi I–IV, respectively.

October 2016, coll. V.V. Abramov. All material is deposited in ZISP.

Description. *Male* (range for 4 specimens). Body including rostral extension of gnathosoma: 425–520 long, 410–480 wide. *Gnathosoma* 165–170 long and 175–190 wide. Palpal femur: 100–130 long; setae *dF* and *v''F* serrate, 60–75 long, 10–12 wide, seta *v'F* smooth filiform. Palpal genu: setae

dG and *l''G* serrate, 70–85 long and 10–12 wide. Palpal tibia: claw with 10–12 basal teeth, setae *dT*, *l'T*, *l''T* smooth filiform. Palpal tarsus: eupathidium *sul* with 14–15 tines, eupathidium *acm* with 2–5 fine basal tines, solenidium ω small stick-like, setae *u'*, *u''* smooth filiform (Fig. 2B). Peritremes shaped as wide arch with small acute apex (roughly as onion dome), each with 8 chambers. Rostral shield

weakly ornamented with fine punctures. *Idiosoma* 300–355 long. Propodonal shield 150–190 long, with 8 pairs of setae (Fig. 1). All propodonal setae narrowly lanceolate and serrate, 10–12 wide; length of setae: *vi* 55–100, *ve* 80–105, *si* 55–100, *se* 75–105, *c1* 30–55, *c2* 35–50, *c3* 30–50, and *c4* 45–80. Hysterosomal shield 125–140 long, with 6 pairs or lanceolate serrate setae 10–12 wide; length of these setae: *d1* 30–35, *d2* 30–55, *e2* 30–50, *e1* 30–50, *f1* 20–30, and *f2* 35–40. Setae *e1* inserted posterior to level of setae *e2*. Pseudanal setae *ps1*–*ps3* present, narrowly lanceolate; seta *ps2* with subapical tine. Genital setae *g1* and *g2* rod-like, about 5 and 7 long, respectively. 2 pairs of filiform agential setae *ag* present. Aedeagus about 20–28 long.

Tarsi I–IV with solenidion $\omega 1$; length and position of this solenidia: on tarsus I 90–105, situated dorsally; on tarsus II 40–45 long, situated ventrally; on tarsus III 30–40, situated ventrally, on tarsus IV 50–60, situated ventrally (Fig. 2C–F). Claws of all tarsi without basal angle. Solenidia ϕ of tibia I and II shaped as short sticks, about 5 long; solenidion $\sigma 1$ of genu I rudimentary, shaped as minute button. Length of legs I 405–475; length of tarsi: I—100–130, II—75–95, III—75–100, IV—90–105.

Remarks. Among previously described species for which males are known, *Mexecheles virginensis* is most similar to *M. thailandensis* and *M. votandinii* (Jeffrey 1975; Faungarworn and Lekprayoon 2010). In males of these species, all dorsal idiosomal setae are narrowly lanceolate or spatuliform and densely serrate. In *M. shiva*, these setae are rod-like and sparsely serrate. Males of *M. virginensis* are distinguished by the form of their peritremes; they form an acute anterior angle, i.e., they are shaped like an onion dome (vs. trapezoid-shaped in *M. thailandensis* and rounded in *M. votandinii*). It is necessary to note that male specimens of *M. virginensis* described herein are noticeably smaller than those described from India (Fain and Bochkov 2001). The body length of European specimens is 425–520 (vs. 570 in the Indian specimen) and the gnatasoma is punctured dorsally (vs. ornamented with short irregular dashes).

Distribution. This species was originally described from the USA, Virginia (Baker 1949). Later on, it was recorded in a number of localities of the New World: USA (Alabama, Louisiana, Mississippi, Nevada and Utah), Canada and Honduras (Smiley and Moser 1970; Summers and Price 1970). In the Old World, it was recorded in Finland

(Penttinen *et al.* 2013), Russia (Tula Region) (Bochkov and Abramov 2016) and India (Kashmir) (Fain and Bochkov 2001).

ACKNOWLEDGEMENTS

The authors express their gratitude to late Dr. A.V. Bochkov (Zoological Institute RAS, St. Petersburg, Russia) for his help with the description technique.

REFERENCES

- Baker, E.W. 1949. A review of the mites of the family Cheyletidae in the United States National Museum. *Proceedings of the United States National Museum*, 99: 267–320.
- Bochkov, A. V. 2004. [Mites of the family Cheyletidae (Acari: Prostigmata); phylogeny, distribution, evolution and analysis of host-parasite relationships]. *Parazitologiya*, 38: 122–138. [In Russian]
- Bochkov, A.V. 2009. A review of mites of the parvorder Eleutherengona (Acariformes: Prostigmata)—permanent parasites of mammals. *Acarina*, Supplement: 1–149.
- Bochkov, A.V. and Abramov, V.V. 2016. To fauna of the free-living Cheyletidae (Acariformes: Cheyletoidea) of the European part of Russia. *Systematic and Applied Acarology*, 21 (3): 335–346.
- Bochkov, A.V. and Fain, A. 2001. Phylogeny and system of the Cheyletidae (Acari: Prostigmata) with special reference to their host-parasite associations. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 71: 5–36.
- Bochkov, A.V. and Ochoa, R. 2005. Three new cheyletid species (Acari: Cheyletidae) from the National Museum of Natural History Entomological Collection. *Acarina*, 12 (2): 137–144.
- De Leon, D. 1962. Three new genera and seven new species of cheyletids (Acarina: Cheyletidae). *Florida Entomologist*, 45 (3): 129–137.
- Fain, A. and Bochkov, A.V. 2001. A review of some genera of Cheyletid mites (Acari: Prostigmata) with descriptions of new species. *Acarina*, 9 (1): 47–95.
- Faungarworn, M. and Lekprayoon, C. 2010. Two new species of cheyletid mites (Acari: Prostigmata) from Thailand. *Zootaxa*, 2494: 59–68.
- Grandjean, F. 1939. Les segments postlarvaires de l'hysterosoma chez les oribates (Acarines). *Bulletin de la Société Zoologique de France*, 64: 273–284.
- Grandjean, F. 1944. Observations sur les acaries de la famille des Stigmaeidae. *Archives des Sciences Physiques et Naturelles*, 26: 103–131.
- Jeffrey, I.G. 1975. An undescribed species of *Mexecheles* (Acarina: Cheyletidae) from a Scottish farm. *Acarologia*, 17 (4): 668–670.

- Kethley, J. 1990. Acarina: Prostigmata (Actinedida). *In*: D.L. Dindal (Ed.). *Soil Biology Guide*. Wiley, New York, 667–756.
- Penttinen, R., Viiri, H. and Moser, J. C. 2013. The mites (Acari) associated with bark beetles in the Koli national park in Finland. *Acarologia*, 53: 3–15.
- Smiley, R. L. and Mooser, J. C. 1970. Three cheyletids found with pine bark beetles (Acarina: Cheyletidae). *Proceedings of the Entomological Society of Washington*, 72: 229–236.
- Summers, F. M. and Price, D. W. 1970. Review of the mite family Cheyletidae. *University of California Publications in Entomology*, 61: 1–153.
- Volgin, V.I. 1955. [The family Cheyletidae Leach—predaceous mites]. *In*: E.N. Pavlovsky (Ed.). Kleshchi Gryzunov Fauny SSSR. Opredeliteli po Faune SSSR, 59. Academy of Sciences of the USSR, Moscow–Leningrad, pp. 152–176. [In Russian]
- Volgin, V. I. 1969. [Mites of the family Cheyletidae of the World]. *In*: Opredeliteli po faune SSSR izdavaemye Zoologicheskim Institutom Akademii Nauk SSSR, 101. Nauka, Leningrad, 432 pp. [In Russian]
- Zhang, Z.-Q., Fan, Q.-H., Pesic, V., Smith, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T., Amrine, J.W., Beron, P., Lin, J., Gabrys, G., and Husband, R. 2011. Order Trombidiformes Reuter, 1909. *In*: Z.-Q. Zhang (Ed.). *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148: 129–138.