

A NEW SPECIES OF *EUSTIGMAEUS* BERLESE (ACARI: PROSTIGMATA: STIGMAEIDAE) FROM NORTHWESTERN IRAN

**K. Haddad Irani-nejad¹, P. Lotfollahi¹, A. Akbari¹, M. Bagheri²
and E. A. Ueckermann^{3, 4}**

¹Department of Plant Protection, Faculty of Agriculture, University of Tabriz, Tabriz, Iran;
e-mails: Khaddad@tabrizu.ac.ir; Prslotollahy@yahoo.com

²Department of Plant Protection, Faculty of Agriculture, University of Maragheh, Maragheh, Iran;
e-mail: mbagheri20022002@yahoo.com

³ARC-PPRI, Private Bagx134, Queenswood, Pretoria 0121, South Africa; e-mail: UeckermannE@arc.agric.za

⁴School of Environmental Sciences and Development, North-West University, Potchefstroom Campus 2520, South Africa

ABSTRACT: A new species of *Eustigmaeus* Berlese (Acari: Prostigmata: Stigmeidae), *Eustigmaeus azerbaijanensis* sp. n. is described and illustrated; The new species differs from the closely related species, *Eustigmaeus coronaria* Kuznetzov, 1977, in the shape of dorsal, ventral and coxal setae and number of paragenital setae. A key to the Iranian species of *Eustigmaeus* is provided.

KEY WORDS: Acari, Stigmeidae, *Eustigmaeus*, new species, Iran

INTRODUCTION

Eustigmaeus Berlese is one of the main genera of the Stigmeidae and contains about 100 species (Dogan et al. 2003, Faraji et al. 2007). Members of the genus *Eustigmaeus* live on mosses, lichens, grass, litter and ant nests or they are bryophagous (Summers and Price 1961; Chaudhri 1965; Wood 1971, 1972, 1973; Gerson 1972a, b; Tseng 1982; Dogan et al. 2003). To date six species of the genus *Eustigmaeus* have been reported from Iran, namely: *Eustigmaeus nasrinae* Khanjani et Ueckermann, 2002, *E. segnis* (Koch, 1836), *E. jiangxiensis* Hu, Chen et Huang, 1996, *E. ornatus* Ueckermann et Meyer, 1987, *E. spathatus* Ueckermann et Meyer, 1987, and *Eustigmaeus sculptus* Doğan, Ayyıldız et Fan, 2003. In this paper, a new species, *Eustigmaeus azerbaijanensis* sp. n. from Iran is described and illustrated. A key to the females of all known species of *Eustigmaeus* from Iran is given.

MATERIAL AND METHODS

Stigmeid mite fauna of East Azerbaijan Province, Iran was sampled in 2009. The new species, *Eustigmaeus azerbaijanensis* sp. n. was collected from an apple orchard at Shendabad (Shabestar) on 15 August 2009 by A.Akbari. Specimens were transferred to the acarology laboratory of the Plant Protection Department, Faculty of Agriculture, University of Tabriz and the mites extracted using a Berlese funnel or direct observation under a stereo microscope. Mites were cleared with Nesbitt's and mounted in Hoyer's medium. Identifications were done with a phase contrast microscope (Olympus, BX41). Illustrations were made using a drawing tube. The leg and idiosomal chaetotaxy

of the species description follows Grandjean (1939, 1944) as adapted by Kethley (1990). All measurements are given in micrometers (μm).

Key to the Iranian species of *Eustigmaeus* Berlese

1. Eyes absent *E. azerbaijanensis* sp. n.
- Eyes present 2
2. Setae c_1 and d_1 longer than 50 μm 3
- Setae c_1 and d_1 shorter than 50 μm 4
3. Coxisternal shields smooth and divided. *E. ornatus* Ueckermann et Meyer, 1987
- Coxisternal shields reticulated and fused *E. segnis* (Koch, 1836)
4. Femur II with five setae 5
- Femur II with 4 setae 6
5. Three pairs of ag setae *E. sculptus* Dogan, Ayyıldız et Fan, 2003
- Two pairs of ag setae *E. jiangxiensis* Hu, Chen et Huang, 1996
6. Dorsal body setae short and spatulate *E. spathatus* Ueckermann et Meyer, 1987
- Dorsal body setae densely setose, most terminating as blunt tip *E. nasrinae* Khanjani et Ueckermann, 2002

Stigmeidae Oudemans, 1931

Type genus: *Stigmeus* Koch, 1836

***Eustigmaeus* Berlese, 1910**

Eustigmaeus Berlese, 1910: 206 (= *Ledermuelleria* Oudemans, 1923)

Type species: *Stigmeus kermesinus* Koch, 1841

Diagnosis. Idiosoma globular and red to straw colored. Gnathosoma partly covered by overhang-

ing prodorsum; Chelicerae separate; palp 5-segmented; palptibial claw well developed, with small accessory claw at its base; terminal eupathidia on palptarsus basally fused and split into 3 long prongs; subcapitulum with 2 pairs of subcapitular setae, *m* and *n*. Dorsum covered with 3 unpaired shields mostly ornamented. Prodorsal shield typically bearing 4 pairs of setae (*vi*, *ve*, *sci* and *sce*) and with or without 1 pair of eyes; opisthosomal shield bearing 6 pairs of setae and suranal shield (usually situated ventrally) bearing 2 pairs of setae. Humeral shields triangular, displaced to a ventral or ventrolateral position bearing *c*₂. Coxisternal shields I-II and III-IV present fused or divided; Aggenital shield with 1 to 3 pairs of setae. Pseudanal shield bearing 3 pairs of setae. Legs stout, shorter than idiosoma; tarsal claws robust; empodium with 3 pairs of tenant hairs before extending beyond tips of claws. The genus *Eustigmaeus* includes about 100 species distributed worldwide (Faraji et al. 2007).

Eustigmaeus azerbaijanensis

Haddad, Lotfollahi et Akbari sp. n.

Figs. 1-8

Female (n=13). Relatively large species, robust and sclerotized; idiosoma globular. Measurements of holotype (variations in measurements of paratypes in parentheses): Length of idiosoma 379 (348–420), width 308 (291–376) (Fig. 1).

Gnathosoma. Palpal setation: tarsus with 1 terminal tridentate eupathidium + 1 solenidion + 5 tactile setae; tibia with 1 well-developed claw + 1 accessory claw + 2 tactile setae; genua with 2 tactile setae; femur with 3 serrate setae (Fig. 3). Accessory claw spine-like. Subcapitular setae *m* longer than *n*, *m* = 35 (32–35), *n* = 28(22–28); *m-m* = 35 (32–35), *n-n* = 35 (30–35), *m-n* = 16 (13–16) (Fig. 2).

Dorsum (Fig. 1). Covered with 3 unpaired shields ornamented with distinct polygonal cells, not vacuolated. Prodorsal shield bearing 4 pairs of setae (*vi*, *ve*, *sci* and *sce*), without eyes and *pobs*; ratios: *vi*: *vi-vi* = 3.6 (3.6–4.1), *ve*: *sci* = 1.2, *sce*: *sci* = 1.1; setae *vi* 127(125–142), *ve* 166 (166–174), *sci* 142 (142–150), *sce* 155 (150–155); distances: *vi-vi* 35, *vi-ve* 43 (41–47), *ve-ve* 85 (79–87), *ve-sci* 43 (43–60), *sci-sci* 163 (163–183), *sci-sce* 51 (44–51). Opisthosomal shield with 6 pairs of setae. Humeral shields triangular, displaced to a ventrolateral position and bears *c*₂; setal lengths: *c*₁ 169 (158–169), *c*₂ 126 (120–130), *d*₁ 177 (166–177), *d*₂ 166 (158–166), *e*₁ 174 (158–178), *e*₂ 182 (178–185), *f*₁ 148 (148–150); ratios

*c*₁: *c*₁-*c*₁ = 1.6, *e*₁: *e*₁-*e*₁ = 1.7 (1.4–1.7), *c*₁-*c*₁: *d*₁-*d*₁: *e*₁-*e*₁: *f*₁-*f*₁ = 1.4–1.5: 1.3–1.4: 1.5–1.6: 1.1–1.2; distances: *c*₁-*c*₁ 103 (99–103), *c*₁-*d*₁ 87 (85–87), *d*₁-*d*₁ 99 (92–99), *d*₁-*d*₂ 74 (74–87), *d*₂-*d*₂ 237 (237–253), *d*₁-*e*₁ 92 (92–101), *e*₁-*e*₁ 103 (103–111), *e*₁-*e*₂ 63 (63–68), *e*₂-*e*₂ 221 (213–223), *e*₁-*f*₁ 55 (44–57), *f*₁-*f*₁ 84 (79–84). Suranal shield (situated ventrally) reticulated and bearing 2 pairs of setae, *h*₁ 95 (95–103), *h*₂ 54 (54–62). Dorsal setae as figured (Fig. 8).

Venter (Fig. 2). Covered with striae except for coxal and anogenital areas. Ventral setae *1a*, *3a* and *4a* equal in length, ratio *1a*: *3a*: *4a* = 1.3–1.6: 1.3–1.6: 1.2–1.6; lengths: *1a* 32 (27–32), *3a* 32 (26–32) and *4a* 32 (24–32). Aggenital area with 2 pairs of setae, *ag*₁ 20 (14–20), *ag*₂ 24 (22–24); 3 pairs of pseudoanal setae, lengths: *ps*₁ 27 (18–28), *ps*₂ 32 (17–32), *ps*₃ 38 (35–38).

Legs (Figs. 4–7). Length: leg I 240 (240–257), leg II 225 (221–225), leg III 234 (161–234), leg IV 266 (245–266). Counts of setae (solenidia and setae *k* not included) on legs I–IV: coxae 2, 2, 2, 2; trochanters 1, 1, 2, 1; femora 6, 5, 3, 2; genua 4, 3+1*k*, 1, 1; tibiae 5+1*φ*+1*φp*, 5+1*φp*, 5+1*φp*, 5+1*φp*; tarsi 13+1*ω*, 9+1*ω*, 7+1*ω*, 7+1*ω*. Lengths of solenidia: *Iω* 17, *IIω* 17, *IIIω* 15, *IVω* 6.

Male. Unknown.

Type materials. Holotype female and 13 female paratypes: Shendabad, East Azerbaijan, Iran, from soil in an apple orchard, 15 August 2009, coll. A. Akbari.


Type material deposition. Holotype and 10 paratypes are deposited in the Acarological Collection, Department of Plant Protection, Faculty of Agriculture, University of Tabriz, Tabriz, Iran. Three paratypes will be deposited in the Arachnidida Collection of Plant Protection Research Institute, Pretoria, South Africa.

Remarks. The new species resembles *Eustigmaeus coronaria* Kuznetzov, 1977 in the body shape, arrangement and ornamentation of shields and number of setae on all segments of legs, however, it differs from this species in that dorsal setae are not inflated basally or sheathed, ventral and coxal setae are not serrated and by the presence of only two pairs of paragenital setae (vs. 3 pairs in *E. coronaria*).

Etymology. This species is named after the Province where it was collected.

ACKNOWLEDGEMENTS

This project was supported by the research division of the University of Tabriz, Iran, which is greatly appreciated.



Figs 1–8. *Eustigmaeus azerbaijanensis* sp. n., female: 1 — dorsal view; 2 — ventral view; 3 — palp; 4 — leg I (ω : solenidion on the tarsus; ϕ : solenidion on anterior of tibia; ϕp : solenidion on posterior of tibia); 5 — leg II; 6 — leg III; 7 — leg IV; 8 — dorsal seta.

REFERENCES

- Chaudhri, W.M. 1965. New mites of the genus *Ledermuelleria*. *Acarologia*, 7: 467–486.
- Dogan, S., Ayyildiz, N. and Fan, Q.-H. 2003. Descriptions of two new species and a newly recorded species of *Eustigmaeus* from Turkey (Acari: Stigmeidae). *Systematic and Applied Acarology*, 8: 131–144.
- Faraji, F., Ueckermann, E.A. and Bakker, F. 2007. First record of *Eustigmaeus jiangxiensis* Hu, Chen and Huang (Acari: Stigmeidae) from France with a key to the european species of *Eustigmaeus* Berlese, 1910. *International Journal of Acarology*, 33 (2): 145–151.
- Gerson, U. 1972a. A Redescription of *Ledermuelleria frigida* Habeeb (Acarina: Prostigmata: Stigmeidae). *Acta Arachnologica*, 24: 15–28.
- Gerson, U. 1972b. Mites of the genus *Ledermuelleria* (Prostigmata, Stigmeidae) associated with mosses in Canada, *Acarologia*, 13: 319–343.
- Grandjean, F. 1939. Les segments postlarvaires de l'hysterosoma chez les oribates (Acariens). *Bulletin Societe Zoologie France*, 64: 273–284.

- Grandjean, F. 1944. Observations sur les Acariens de la famille des Stigmeidae. *Archives des Sciences physiques et naturelles*, 26: 103–131.
- Kethley, J. 1990. Acarina: Prostigmata (Actinedida). In: D.L. Dindal (Ed.). *Soil Biology Guide*. John Wiley & Sons, New York, pp. 667–756.
- Summers, F.M. and Price, D.W. 1961. New and redescription species of *Ledermuelleria* from North America (Acarina: Stigmeidae). *Hilgardia*, 31: 369–387.
- Tseng, Y.H. 1982. Mites of the family Stigmeidae of Taiwan with key to genera of the world (Acarina: Prostigmata). *Phytopathologist and Entomologist, National Taiwan University*, 9: 1–52.
- Wood, T.G. 1971. Stigmeidae (Acari: Prostigmata) from the British Solomon Islands. *Acarologia*, 13: 65–87.
- Wood, T.G. 1972. New and redescribed species of *Ledermuelleria* Oudms and *Villersia* Oudms (Acari: Stigmeidae) from Canada. *Acarologia*, 13: 301–318.
- Wood, T.G. 1973. Revision of Stigmeidae (Acari: Prostigmata) in the Berlese Collection. *Acarologia*, 15: 76–95.