REDEFINITION OF THE GENUS *CERATTOMA* (ACARI: HETEROSTIGMATA: PYGMEPHORIDAE) WITH REDESCRIPTION OF *C. URSULAE* FROM EUROPEAN RUSSIA

Alexander A. Khaustov^{1*}, Vladimir V. Abramov² and Andrey N. Bobylev¹

¹Institute of Environmental and Agricultural Biology (X-BIO), Tyumen State University,
Tyumen, Russia

²Independent Researcher, Suворov, Tula Region, Russia

*corresponding author; e-mail: alkhaustov@mail.ru

ABSTRACT: The genus *Cerattoma* Mahunka, 1972 (Acari: Pygmephoridae) is redefined. *Cerattoma ursulae* (Krczal, 1959) is redescribed, based on phoretic females collected under the elytra of a *Crypticus* sp. (Coleoptera: Tenebrionidae) darkling beetle in European Russia. An updated key to the genera of the Pygmephoridae family is provided.

KEY WORDS: Heterostigmatina, Pygmephoridea, systematics, morphology, phoresy.

DOI: 10.21684/0132-8077-2019-27-1-53-63

INTRODUCTION

The cosmopolitan family Pygmephoridae, the second largest in the Pygmephoridea superfamily, includes 31 genera and more than 300 species (Khaustov and Trach 2018). In all likelihood, all pygmephorid mites are fungivorous (Kaliszewski *et al.* 1995).

The genus name *Cerattoma* Mahunka was first mentioned in Mahunka (1970) as created in 1969. However, the description of the genus *Cerattoma* was not published and type species was not designated in 1969 (Mahunka 1972). The brief description and type designation (type species *Pygmephorus ursulae* Krczal, 1959) of *Cerattoma* were provided in Mahunka (1972). He placed five species in the genus *Cerattoma*: *C. ursulae* (Krczal, 1959), *P. ceratophyi* (Krczal, 1959), *C. endroedyoungai* (Mahunka, 1965), *C. szekessyi* Mahunka, 1970, and *C. graeca* Mahunka, 1972. Cross and Moser (1971) created the subgenus *Pygmephorellus* of the genus *Pygmephorus* Kramer, 1877 and placed five species in it: *P. ceratophyi* (Krczal, 1959) (type species), *P. ursulae* (Krczal, 1959), *P. karafati* (Krczal, 1959), *P. brachycerus* Cross and Moser, 1971, and *P. bennetti* Cross and Moser, 1971. Cross (1974) considered *Cerattoma* sensu Mahunka (1970) (with type species *P. ceratophyi*) as an invalid generic name and replaced it by *Pygmephorellus* and moved *P. karafati* and *P. bennetti* to *Elattoma* Mahunka, 1969. Apparently, Cross (1974) did not know about the publication of Mahunka (1972) where he designated *P. ursulae* as type species. Sevastianov (1978) also used the generic name *Cerattoma* Mahunka, 1969 with the type species *Pygmephorus ceratophyi* Krczal, 1959. It is not clear why Cross (1974) and Sevastianov (1978)

considered *P. ceratophyi* as the type species of *Cerattoma*.

During the study of insect-associated mites, a rare species *Cerattoma ursulae* was discovered in European Russia. This is the first record of this species from Russia. The study of the morphology of this species revealed several apomorphic characters, which are found neither in the type nor in most other *Pygmephorellus* species. Based on these characters, we redefine the genus *Cerattoma* Mahunka, 1972 and redescribe its type species, *C. ursulae*.

The latest key to the Pygmephoridae genera was provided by Khaustov (2015). Since that time, several genera were described or redefined (Rahiminejad *et al.* 2015, 2016; Khaustov *et al.* 2017a, b, 2018). Here, we provide an updated key to the Pygmephoridae genera.

MATERIAL AND METHODS

The mites were collected under the elytra of a single beetle *Crypticus* sp. that was captured by V. Abramov in the Tula Region, Russia. The mites were mounted in Hoyer's medium. The terminology follows that of Lindquist (1986); the nomenclature of subcapitular setae and the designation of cheliceral setae follow those of Grandjean (1944, 1947), respectively. The systematics of Pygmephoridea follows that of Khaustov (2004, 2008). All measurements are given in micrometers (μm). For leg chaetotaxy, the number of solenidia is provided in parentheses. DIC micrographs were taken using the Carl Zeiss Axio Imager A2 compound microscope and the AxioCam 506 color digital camera. For SEM microscopy, alcohol-preserved mites were dried in freeze-drying device JFD 320 (JEOL, Japan), dusted with gold, and scanned with the aid

Fig. 1. *Cerattoma ursulae* (Krczal, 1959), female: A—dorsum of the body; B—venter of the body. Legs omitted.

of a JEOL–JSM-6510LV SEM microscope. All materials have been deposited in the mite collection of the Tyumen State University Museum of Zoology, Tyumen, Russia.

SYSTEMATICS

Family **Pygmephoridae** Cross, 1965

Genus ***Cerattoma*** Mahunka, 1972

Type species: *Pygmephorus ursulae* Krczal, 1959, by original designation.

Diagnosis. Female. Body well sclerotized, oval. Gnathosomal capsule of about equal length and width, dorsally with one pair of cheliceral setae (*cha*), setae *chb* absent; postpalpal setae (*pp*) present; palps prominent, with two pairs of setae (*dFe*, *dGe*); tibial claw large; palpal solenidion (*sol*)

well developed, accessory setigenous structure (*ass*) large, mushroom-like. Palp tibiotarsus with tiny distal eupathidium. Subcapitular setae (*m*) present. Subcapitulum with one pair of oval alveolar pits *n* in posterior half. Pharyngeal pumps tripartite, situated on long and thin oesophagus; pump 1 bow-shaped, smooth, far separated from pump 2, pumps 2 and 3 subuqual, oval, situated close to each other. Prodorsum with three pairs of setae (*v*₁, *v*₂, *sc*₂), one pair of clavate trichobothria (*sc*₁) and one pair of oval stigmata; prodorsal shield not divided. Tracheal trunks thin, long. Posterior part of prodorsal shield covering anterior part of tergite C. Tergite C with two pairs of setae (*c*₁, *c*₂); tergite D with one pair of setae (*d*) and one pair of round cupules *ia*; tergite EF with two pairs of setae (*e*, *f*), cupules *im* absent; tergite H with two pairs

Fig. 2. *Cerattoma ursulae* (Krczal, 1959), female: A—gnathosoma in dorsal view; B—gnathosoma and pharyngeal pumps in ventral view.

Fig. 3. *Cerattoma ursulae* (Krczal, 1959), female: A—right leg I in dorsal view; B—right leg II in dorsal view.

of setae (h_1, h_2) and one pair of round cupules *ih*. Coxal fields I with three pairs of setae (1a, 1b, 1c), setae 1b not modified; coxal fields II with two pairs of setae (2a, 2c); coxal fields III with three pairs of setae (3a, 3b, 3c); coxal fields IV with three pairs of setae (4a, 4b, 4c). Pseudanal segment with three pairs of setae (ps_{1-3}). Apodemes 1 (ap1) well developed, thick, apodemes 2 (ap2) well developed, joined with well-developed prosternal apodeme (appr); appr joined with sejugal apodeme (apsej); secondary transverse apodeme absent; apodemes 3 (ap3) weak, diffuse; poststernal apodeme (appo) well developed, fused with long apodemes 4 (ap4) and well developed apodemes 5 (ap5). Posterior margin of posterior sternal plate entire. Anterior genital sclerite (ags) small, bell-like, median genital sclerite (mgs) present, posterior genital sclerite (pgs) widened anteriorly and narrowed posteriorly (Fig. 5D). Leg I 4-segmented, with massive tibiotarsus. Tarsal claw large, twisted ventrally and arising from apicomedial portion of tibiotarsus. Unguinal setae fused to form bifurcate structure opposing the tarsal claw and twisted dorsally (Fig.

7D). Tibiotarsus without pinnaculum. Seta *d* of femur I hook-like. Seta *k* smooth, blunt-ended. Legs II–IV each with one pair of claws and empodium with characteristic pointed tip (Fig. 7E). Claws on tarsi II and III thickened basally, on tarsus IV simple. Femora III and IV divided into basi- and telofemur. Leg setation: leg I; Tr 1 (v'), Fe 4 (d, l', l'', v''), Ge 4 (l', l'', v', v''), TiTa 17(4) ($d, l', l'', v', v'', k, tc', tc'', p', p'', ft', ft'', pv', pv'', pl', pl'', s, \omega_1, \omega_2, \phi_1, \phi_2$); leg II: Tr 1 (v'), Fe 3 (d, l', v''), Ge 3 (l', v', l''), Ti 4(1) (d, l', v', v'', ϕ), Ta 6(1) ($tc'', tc'', pl'', pv', pv'', u', \omega$); leg III: Tr 1 (v'), Fe 2 (d, v'), Ge 2 (l', v'), Ti 4(1) (d, l', v', v'', ϕ), Ta 6 ($tc'', tc'', pl'', pv', pv'', u'$); leg IV: Tr 1 (v'), Fe 2 (d, v'), Ge 1 (v'), Ti 4(1) (d, l', v', v'', ϕ), Ta 6 ($pl'', tc', tc'', u', pv', pv''$).

Male and larva unknown.

Species included. The genus *Cerattoma* includes two species: *C. ursulae* (Krczal, 1959) and *C. graeca* Mahunka, 1972.

Distribution and habitat. *C. ursulae* was described from the soil obtained from a potato field in Germany (Krczal 1959). It was also recorded

Fig. 4. *Cerattoma ursulae* (Krczal, 1959), female: A—right leg III in dorsal view; B—right leg IV in dorsal view.

from soils in Hungary (Mahunka 1981) and Ukraine (Sevastianov 1978). This species is phoretic under

the elytra of the *Crypticus* sp. darkling beetle in European Russia (present data). *Cerattoma graeca*

Fig. 5. DIC micrographs of *Cerattoma ursulae* (Krczal, 1959), female: A—prodorsum; B—anterior sternal plate; C—posterior sternal plate; D—genital sclerites.

was described from the leaf litter associated with *Quercus pubescens*, in Greece (Mahunka 1972).

Remarks. We retain the incompletely described species *Cerattoma graeca* in the genus *Cerattoma*, based on its unusual, twisted ventrally tarsal claw on tibiotarsus I, which was not illustrated in the original description, but was well-described in the text (Mahunka 1972). Other characters are also very similar to the type species of the genus *Cerattoma*.

Differential diagnosis. The genus *Cerattoma* is very similar to *Pygmephorus* Cross and Moser, 1971 in the same idiosomal and leg chaetotaxy, as well as in the shape of seta *d* of femur I. It can be distinguished from *Pygmephorus*, however, by an unusual tarsal claw on tibiotarsus I, which is twisted ventrally and arises from the apicomedial portion of tibiotarsus (in the normal dorsal position in *Pygmephorus*). *Cerattoma* can also be distinguished from *Pygmephorus* by the absence of cupules *im* on tergite EF (present in *Pygmephorus*) and by an unusual shape of empodia, with

narrow pointed tips on tarsi II–IV (tips of empodia rounded in *Pygmephorus*).

***Cerattoma ursulae* (Krczal, 1959)**

Pygmephorus ursulae Krczal, 1959, p. 495, figs. 44–45

Cerattoma ursulae: Mahunka 1970, p. 159

Pygmephorus (*Pygmephorus*) *ursulae*; Cross and Moser 1971, p. 55.

Cerattoma ursulae: Mahunka 1972, p. 73

(Figs. 1–7)

Redescription. *Female*. Length of idiosoma 285–325, width 150–180.

Gnathosoma (Figs 2, 6F). Length of gnathosomal capsule 23–27, width 23–28. Dorsal median apodeme present. All gnathosomal setae smooth. Setae *cha* blunt-ended, other gnathosomal setae pointed. Cheliceral levers very small, elongate-oval; cheliceral stylets indistinct. Pharyngeal pump 2 weakly transversely striated, other pumps smooth

Fig. 6. SEM micrographs of *Cerattoma ursulae* (Krczal, 1959), female: A—general view dorsally; B—general view laterally; C—anterior half of prodorsum; D—general view ventrally; E—prosoma in ventral view; F—hypognathous gnathosoma.

(Fig. 2B). Lengths of gnathosomal setae: *cha* 7–9, *pp* 4–5, *dFe* 7–8, *dGe* 12–14, *m* 11–13.

Idiosomal dorsum (Figs. 1A, 5A, 6A, C). All dorsal shields with numerous and rather large dimples (Fig. 6A). Trichobothria distinctly barbed (Fig. 7A). All dorsal setae blunt-ended. Setae *e* and *h*₂ smooth, other dorsal setae weakly barbed. Cupules *ia* on tergite D, and *ih* on tergite H small, round. Setae *e* and *h*₂ situated

distinctly anteriad bases of setae *f* and *h*₁, respectively. Posterior margin of prodorsal shield weakly sclerotized and usually undulate. Tergites D and EF each with a pair of round porous areas located mediad setae *d* and *f*, respectively. Lengths of dorsal setae: *v*₁ 18–21, *v*₂ 17–18, *sc*₂ 39–47, *c*₁ 35–44, *c*₂ 44–52, *d* 36–44, *e* 14–16, *f* 37–41, *h*₁ 45–50, *h*₂ 10–13. Distances between setae: *v*₁–*v*₁ 8–11, *v*₂–*v*₂ 32–38, *sc*₂–*sc*₂ 39–43,

Fig. 7. SEM micrographs of *Cerattoma ursulae* (Krczal, 1959), female: A—lateral part of prodorsum and leg I; B—leg I in dorsal view; C—leg I in ventral view; D—claw and opposing structure (fused unguinal setae) on tibiotarsus I; E—tip of tarsus II in ventral view; F—tibia II in dorsal view.

c_1-c_1 46–54, c_1-c_2 27–32, $d-d$ 88–100, $e-f$ 9–11, $f-f$ 84–99, h_1-h_1 69–82, h_1-h_2 9–11.

Idiosomal venter (Figs 1B, 5B–D, 6D, E). Ventral plates with numerous rather large dimples (Figs 5B, C). Setae $1b$, $1c$ and ps_3 weakly barbed, other ventral setae smooth. Setae $1b$, $1c$, $4b$ and ps_{1-3} pointed, other ventral setae weakly blunt-ended. Aggenital plate with a pair of well developed alveolar pits located posterolaterad setae $4b$. Lengths

of ventral setae: $1a$ 15–16, $1b$ 22–24, $1c$ 19–21, $2a$ 16–17, $2c$ 15–17, $3a$ 17–18, $3b$ 13–16, $3c$ 17–19, $4a$ 12–13, $4b$ 23–27, $4c$ 16–17, ps_1 10–12, ps_2 9–10, ps_3 21–24.

Legs (Figs. 3, 7). Leg I (Figs. 3A, 7A–D). Tibiotarsus slightly thickened. Lengths of solenidia ω_1 17–18, ω_2 21–24, ϕ_1 10–11, ϕ_2 10–11; solenidion ϕ_1 clavate, other solenidia baculiform. Seta l' of femur and k of tibiotarsus blunt-ended and

smooth, other leg setae (except eupathidia) pointed and barbed. Leg II (Figs. 3B, 7E, F). Solenidion ω 17–18 finger-shaped, solenidion ϕ 4–5 weakly clavate, situated in depression (Fig. 7F). Setae l' of femur, v' of genu and u' of tarsus weakly blunt-ended, other setae pointed. Seta l' of femur smooth, other setae weakly barbed. Leg III (Fig. 4A). Solenidion ϕ 3 weakly clavate, situated in depression. Setae v' of trochanter, v' of femur, l' of genu, and u' of tarsus weakly blunt-ended, other setae pointed. All leg setae barbed. Leg IV (Fig. 4B). Solenidion ϕ very small 2–3, located in deep depression and only its tip visible in pore-like opening. All setae barbed. Setae v' of trochanter, v'' of tibia, and pl'' of tarsus blunt-ended, other leg setae pointed.

Material examined. 16 females, Tula Region, vicinity of Suvorov town, under elytra of a single specimen of darkling beetle *Crypticus* sp., collected in pitfall trap, 54°07'N, 36°30'E, September 28, 2018, coll. V.V. Abramov.

Remarks. *Cerattoma ursulae* is very similar to the incompletely described *C. graeca* Mahunka, 1972, but can be distinguished from the latter by the ~1.2 ratio of setae f to h_1 (h_1 is almost two times longer than f in *C. graeca*).

Key to genera of Pygmephoridae

(based on females, after Khaustov 2015, with modifications)

1. Legs I 5-segmented (tibia and tarsus separated)... 2
— Legs I 4-segmented (tibia and tarsus fused).....10
2. Setae e absent.....3
— Setae e present.....6
3. Prodorsum with one or three pairs of simple setae (excluding trichobothria) 4
— Prodorsum with two pairs of simple setae (excluding trichobothria).....*Parasiteroptes* Livshits, Mitrofanov and Sharonov, 1986
4. Prodorsum with three pairs of simple setae (excluding trichobothria).....5
— Prodorsum with one pair of simple setae (excluding trichobothria).....*Pediculitopsis* Mahunka, 1970 (non-phoretic female)
5. Setae $4a$ present, $4c$ absent, body oval
.....*Krczaldania* Sasa, 1961
— Setae $4a$ absent, $4c$ present, body fusiform.....
.....*Siteroptes* Amerling, 1861
6. Setae h_2 present, body oval7
— Setae h_2 absent, body very long, fusiform.....
.....*Metasiteroptes* Cross, 1965

7. Prodorsum with three pairs of simple setae (excluding trichobothria).....8
— Prodorsum with two pairs of simple setae (excluding trichobothria).....*Ultrasiteroptes* Livshits, Mitrofanov and Sharonov, 1986
8. Cupules *im* present, postpalpal setae present, setae $1b$ usually bifurcate, coxal fields II with 2 pairs of setae9
— Cupules *im* absent, postpalpal setae absent, setae $1b$ not bifurcate, coxal fields II usually with 3 pairs of setae*Pediculaster* Vitzthum, 1931 (non-phoretic female)
9. Coxal fields I with two pairs of setae.....*Sevastianoviella* Livshits, Mitrofanov and Sharonov, 1986
— Coxal fields I with three pairs of setae.....*Neositeroptes* Livshits, Mitrofanov and Sharonov, 1986
10. Coxal fields II with three pairs of setae 11
— Coxal fields II with two pairs of setae 16
11. Setae v' of femur and pl'' of tarsus of leg IV not sword-like, claw of tibiotarsus I usually not very large, not striated.....12
— Setae v' of femur and pl'' of tarsus of leg IV sword-like, claw of tibiotarsus I very large, striated, associated with small mammals.....
.....*Pygmephorus* Kramer, 1877
12. Stigmata usually oval or round, sometimes with chambers, not long and narrow 13
— Stigmata long and narrow
.....*Luciaphorus* Mahunka, 1981
13. Tibiotarsus I with claw 14
— Tibiotarsus I without claw.....
.....*Microdispodides* Vitzthum, 1914.
14. Setae l'' of femur I not modified and not similar in shape with d of femur I.....15
— Setae l'' of femur I spine-like, similar in shape with d of femur I *Mahunkania* Rack, 1972.
15. Setae d of femur I spatulate, longer than width of femur I*Pediculaster* Vitzthum, 1931 (phoretic female)(part)
— Setae d of femur I blade-like, shorter than width of femur I *Propygmephorus* Cross, 1974
16. Coxal fields I with two pairs of setae.....17
— Coxal fields I with one or three pairs of setae.... 21
17. Prodorsum with three pairs of simple setae (excluding trichobothria) 18
— Prodorsum with two pairs of simple setae (excluding trichobothria).....*Sasadania* Kurosa, 1989
18. Setae d of femur I not modified 19
— Setae d of femur I spatulate20
19. Trichobothria present
.....*Dudichiana* Mahunka, 1970
— Trichobothria absent*Asensilla* Rack, 1974

20. Tarsus II with four modified, spine-like setae.
..... *Pygmephoroides* Mahunka and Fain, 1989.
— Tarsus II without spine-like setae.....
..... *Pediculaster* Vitzthum, 1931
(phoretic female) (part)
21. Coxal fields I with three pairs of setae..... 22
— Coxal fields I with one pair of setae.....
..... *Geotrupophorus* Mahunka, 1970
22. Tarsi II and III with six setae each, trichobothria present..... 23
— Tarsi II and III with only four setae each (*tc* ' and *pl* ' absent), trichobothria absent.....
..... *Micropygmephorus* Khaustov, Hugo-Coetzee and Ermilov, 2017
23. Setae *d* of femur I longer than width of femur I, spatulate..... 24
— Setae *d* of femur I shorter than width of femur I, spatulate or hook-like 28
24. Prodorsum with three pairs of simple setae (excluding trichobothria) 26
— Prodorsum with two pairs of simple setae (excluding trichobothria)..... 25
25. Pseudanal segment distinctly narrowed and elongate posteriorly..... *Pediculitopsis* Mahunka, 1970 (phoretic form)
— Pseudanal segment of normal shape, evenly rounded posteriorly.....
..... *Apediculaster* Rahiminejad and Hajiqanbar, 2016
26. Setae *h*₂ present, body oval 27
— Setae *h*₂ absent, body fusiform.....
..... *Brasilopsis* Mahunka, 1975 (probably phoretic female of *Metasiteroptes*)
27. Leg IV distinctly shorter than leg III, tarsus IV short, with 3 very long, whip-like setae.....
..... *Acarothorectes* Cross, 1965
— Leg IV not shorter than leg III, tarsus IV not short, sometimes with only one whip-like setae *tc* '.....
..... *Pediculaster* Vitzthum, 1931 (phoretic female)(part)
28. Empodia on tarsi II–IV present..... 29
— Empodia on tarsi II–IV absent *Mesopotamiophorus* Sevastianov and Zahida Al Douri, 1991
29. Genu I with one or two setae, genu IV without setae..... 30
— Genu I with three or four setae, genu IV with one setae..... 31
30. Tibiotarsus I with two solenidia, setae *d* of femur I hook-like, two pairs of pseudanal setae, setae 4*a* and 4*c* absent *Elattoma* Mahunka, 1969
— Tibiotarsus I with four solenidia, setae *d* of femur I spatulate, three pairs of pseudanal setae, setae 4*a* and 4*c* present *Spatulaphorus* Rack, 1993
31. Lateral surfaces of tibiotarsus I without slit-like structures, most setae of anterior and posterior sternal plates not modified 32
— Lateral surfaces of tibiotarsus I with slit-like structures, most setae of anterior and posterior sternal plates spine-like..... *Strephocheir* Mahunka, 1983
32. Stigmata oval, ungual setae (*u*) of tibiotarsus I present as modified structure opposing to tarsal claw 33
— Stigmata long and narrow, ungual setae (*u*) of tibiotarsus I absent
..... *Parapediculaster* Khaustov, 2015
33. Subcapitular setae *m* present, setae 2*b* not bifurcate 34
— Subcapitular setae *m* absent, setae 2*b* bifurcate ...
..... *Metapygmephorus* Rahiminejad, Hajiqanbar and Khaustov, 2015
34. Genu I with four setae, genu II with three setae..... 35
— Genu I with three setae, genu II with two setae..... *Pseudopygmephorus* Khaustov, 2008
35. Cupules *im* present, empodia on tarsi II–III rounded distally, claw on tibiotarsus I situated dorsally..... *Pygmephorus* Cross and Moser, 1971
— Cupules *im* absent, empodia on tarsi II–III pointed distally, claw on tibiotarsus I twisted ventrally *Cerattoma* Mahunka, 1972

ACKNOWLEDGEMENTS

The work of A.A. Khaustov was supported by the Russian Foundation for Basic Research (RFBR), research project № 18-04-01092A.

REFERENCES

- Cross, E.A. 1974. A new genus and two new species of pyemotid mites phoretic upon insects (Acarina: Tarsonemoidea). *Acarologia*, 16 (4): 677–683.
- Cross, E.A. and Moser J.C. 1971. Taxonomy and biology of some Pyemotidae (Acarina: Tarsonemoidea) inhabiting bark beetle galleries in North American conifers. *Acarologia*, 13 (1): 47–64.
- Grandjean, F. 1944. Observations sur les Acariens de la famille des Stigmaeidae. *Archives des Sciences Physiques et Naturelles*, 26: 103–131.
- Grandjean, F. 1947. L'origine pileuse des mors et la chaetotaxie de la mandibule chez les Acariens actinochitineux. *Comptes rendus des Séances de l'Académie des Sciences*, 224: 1251–1254.
- Kaliszewski, M., Athias-Binche, F. and Lindquist, E.E. 1995. Parasitism and parasitoidism in Tarsonemina (Acari: Heterostigmata) and evolutionary considerations. *Advances in Parasitology*, 35: 335–367.

- Khaustov, A.A. 2004. Mites of the family Neopygmephoridae Cross, 1965 stat. n. and their position in Heterostigmata. In: Y.S. Balashov (Ed.), VIII Russian Acarological Conference, St.Petersburg. Zoological Institute of RAS, St.Petersburg, p. 137. [In Russian]
- Khaustov, A.A. 2008. *Mites of the family Scutacaridae of Eastern Palaearctic*. Akademperiodyka, Kiev, 291 pp.
- Khaustov, A.A. 2015. A new genus and species of the family Pygmephoridae (Acari: Pygmephoridae) from southern Chile. *International Journal of Acarology*, 41: 202–209.
- Khaustov, A.A., Çobanoglu, S. and Cilbircioğlu, C. 2017a. Redefinition of the genus *Asensilla* Rack (Acari: Heterostigmata: Pygmephoridae) with redescription of *A. prassei*. *Acarina*, 25 (2): 129–134.
- Khaustov, A.A., Hugo-Coetzee, E.A. and Ermilov S.G. 2017b. A new genus and two new species of Pygmephoridae (Acari: Heterostigmata) associated with *Trinervitermes trinervoides* (Isoptera: Termitidae) from South Africa. *Zootaxa*, 4258 (5): 462–476.
- Khaustov, A.A., Shentu, H. and Ch.-Ch, Ho. 2018. Redefinition of the genus *Luciaphorus* Mahunka, 1981 (Acari: Pygmephoridae) with redescription of *L. perniciosus* Rack, 1983 and *L. auriculariae* Gao, Zou and Jiang, 1990. *Zootaxa*, 4369 (4): 536–556.
- Khaustov, A.A. and Trach, V.A. 2018. New species of *Spatulaphorus* (Acari: Heterostigmata: Pygmephoridae) associated with *Geotrupes baicalicus* (Coleoptera: Geotrupidae) from Altai Republic, Russia. *Systematic and Applied Acarology*, 23: 123–131.
- Krczal, H. 1959. Systematik und Ökologie der Pygmotiden. *Beiträge zur Systematik und Ökologie der mitteleuropäischen Acarina*, 1: 385–625.
- Lindquist, E.E. 1986. The world genera of Tarsonemidae (Acari: Heterostigmata): a morphological, phylogenetic, and systematic revision, with a reclassification of family-group taxa in the Heterostigmata. *Memoirs of Entomological Society of Canada*, 118: 1–517.
- Mahunka, S. 1970. Considerations on the systematics of the Tarsonemina and the description of new European taxa (Acari: Trombidiformes). *Acta Zoologica Academiae Scientiarum Hungaricae*, 16: 137–174.
- Mahunka S. 1972. Neue und interessante Milben aus dem Genfer Museum I. Angaben über die Tarsonemina-Fauna (Acari) aus Kephallinia, Griechenland. *Biologia Gallo-Hellenica*, 4 (1): 71–83.
- Mahunka, S. 1981. The Pygmephoroid fauna of the Hortobágy National Park (Acari: Tarsonemida). *The Fauna of the Hortobágy National Park*, 1: 343–370.
- Rahiminejad, J., Hajiqaanbar, H., Khaustov, A.A. and Talebi, A.A. 2015. A new genus and two new species of the family Pygmephoridae (Acari: Heterostigmata) associated with beetles (Insecta: Coleoptera). *Annales of the Entomological Society of America*, 108: 893–901.
- Rahiminejad, J., Hajiqaanbar, H. and Talebi, A.A. 2016. A new genus and species of the family Pygmephoridae (Acari: Heterostigmata) associated with *Carpelimus rivularis* (Coleoptera: Staphylinidae). *Systematic and Applied Acarology*, 21: 461–470.
- Sevastianov, V.D. 1978. Tarsonemina. In: M.S. Ghilarov (Ed.), *Opredelitel pochvoobitayushchikh kleshchey. Trombidiformes*, Moscow, Nauka, pp. 14–90. [In Russian]